

**UNIVERSITI TEKNOLOGI MARA**

**ANTAGONISTIC EFFECTS  
OF SELECTED MICROBES  
ON *Fusarium* spp.**

**IDA HANI BINTI ALI**

Thesis submitted in fulfilment  
of the requirements for the degree of  
**Master of Science**

**Faculty of Applied Sciences**

**September 2013**

## AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA, Malaysia. It is original and is the result of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any other degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate Universiti Teknologi MARA, Malaysia, regulating the conduct of my study and research.

Name of Student : Ida Hani Binti Ali

Student I.D No. : 2007131439

Programme : Master of Science

Faculty : Faculty of Applied Sciences

Title : Antagonistic Effects of Selected Microbes  
on *Fusarium* spp.

Signature of Student :  .....

Date : September 2013

## ABSTRACT

A study was conducted to evaluate the antagonistic activities of selected fungal and bacterial species which included *Aspergillus nidulans*, *Aspergillus niger*, *Penicillium chrysogenum*, *Trichoderma reesei*, *Bacillus subtilis*, *Pseudomonas aeruginosa* and *Serratia marcescens* towards the plant pathogens, *Fusarium sacchari* and *Fusarium oxysporum* f. sp. *batatas*. All the selected microorganisms were found to have antagonistic activities against the fungal pathogens when evaluated by the dual culture assay on potato dextrose agar at 30°C for duration of 9 days of incubation. The strongest microorganism for both fungal pathogens was *T. reesei* followed by *A. niger*, *P. aeruginosa*, *P. chrysogenum*, *A. nidulans*, *B. subtilis* and *S. marcescens*. The involvement of mycoparasitism or antibiosis activity was also investigated. Microscopic observations of mycelial samples from the interaction area from the dual culture assay plates showed that the hyphae of *Fusarium* spp. were either lysed, fragmented, shriveled, swelled, deformed and became stunted. Deformities, disintegration and lost of the structural integrity of the pathogen conidia were also observed. Antibiosis in the microbial cell-free culture filtrates and methanol crude extracts was detected using agar well diffusion assay and broth assay. Antibiosis activity towards both fungal pathogens was only detected in cell-free culture filtrate and methanol crude extract of *P. aeruginosa* in the broth assay. There were no antibiosis activity showed by the other microbial crude extracts. Analysis of gas chromatography mass spectrometry was detected the presence of compounds namely cycloheptasiloxane, tetradecamethyl and 1,1,1,5,7,7,7-Heptamethyl-3, 3-bis (trimethylsiloxy) tetrasiloxane in the methanol crude extract of *P. aeruginosa* which may responsible to the antibiosis activity. From this study, the selected microbes that have the potential to be biological control agents for plant diseases caused by *F. sacchari* and *F. oxysporum* f. sp. *batatas* are *T. reesei*, *A. niger* and *P. aeruginosa*.

TABLE OF CONTENTS

	Page
AUTHOR’S DECLARATION	ii
ABSTRACT	iii
ACKNOWLEDGEMENTS	iv
TABLE OF CONTENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	x
LIST OF ABBREVIATIONS	xii

CHAPTER ONE: INTRODUCTION

1.1	Background	1
1.2	Problem Statement	2
1.3	Significance of Study	3
1.4	Scope and Limitation	3
1.5	Objectives	4

CHAPTER TWO: LITERATURE REVIEW

2.1	The Genus <i>Fusarium</i>	5
2.2	Taxonomy of <i>Fusarium</i>	5
2.2.1	Toxigenic <i>Fusarium</i> Species	7
2.2.2	<i>Fusarium</i> spp. Associated with Plant Diseases	9
2.2.2.1	<i>F. sacchari</i>	11
2.2.2.2	<i>F. oxysporum</i> f. sp. <i>batatas</i>	13
2.2.2.3	<i>F. oxysporum</i>	15
2.2.2.4	Other <i>Fusarium</i> Species	16
2.3	Microorganisms as Biological Control Agents	18
2.3.1	Fungi	19
2.3.1.1	<i>Aspergillus</i> spp.	19

2.3.1.2	<i>Penicillium spp.</i>	19
2.3.1.3	<i>Trichoderma spp.</i>	20
2.3.2	Bacteria	24
2.3.2.1	<i>Bacillus spp.</i>	24
2.3.2.2	<i>Pseudomonas spp.</i>	24
2.3.2.3	<i>Serratia spp.</i>	24
2.4	Antagonistic Activity by Dual Culture Technique	26
2.5	Mechanisms of Antagonism	28
2.5.1	Competition	28
2.5.2	Mycoparasitism	29
2.5.3	Antibiosis	31

### CHAPTER THREE: MATERIALS AND METHODS

3.1	Materials	33
3.1.1	Microbial Strains	33
3.1.2	Media	34
3.1.3	Preparation of Cycloheximide	35
3.2	Methods	35
3.2.1	Determination of Antagonistic Activity	37
3.2.2	Determination of Mechanisms of Antagonism	39
3.2.2.1	<i>Mycoparasitism Activity</i>	39
3.2.2.2	<i>Detection of Antibiosis Activity</i>	42

### CHAPTER FOUR: RESULTS

4.1	Antagonistic Activity by Dual Culture Assay	49
4.2	Characterization of Mechanisms of Antagonism	56
4.2.1	Mycoparasitism	56
4.2.1.1	<i>Sample Analysis by Light Microscope</i>	56
4.2.1.2	<i>Sample Analysis by Scanning Electron Microscope</i>	60
4.2.2	Detection of Antibiosis Activity	63
4.2.2.1	<i>Agar Well Diffusion Assay Using Antagonist</i>	
	<i>Cell-Free Culture Filtrates</i>	63